

# Copositivity of a class of fourth order symmetric tensors \*

Yisheng Song<sup>†</sup>, Jinjie Liu<sup>‡</sup>

## Abstract

In this paper, we mainly discuss the analytic conditions of copositivity of a class of 4th order 3-dimensional symmetric tensors. For a 4th order 3-dimensional symmetric tensor with its entries 1 or  $-1$ , an analytic necessary and sufficient condition is given for its strict copositivity with the help of the properties of strictly semi-positive tensors. And by means of usual maxi-min theory, a necessary and sufficient condition is established for copositivity of such a tensor also. Applying these conclusions to a general 4th order 3-dimensional symmetric tensor, the analytic conditions are successfully obtained for verifying the (strict) copositivity, and these conditions can be very easily parsed and validated.

**Keywords:** Copositivity, Fourth order tensors, Homogeneous polynomial.

**Mathematics Subject Classification:** 90C23, 15A69, 15A72, 90C30, 90C47

## 1 Introduction

One of the most direct applications of 4th order copositive tensors is to verify the vacuum stability of the Higgs scalar potential model

---

\*The first author's work was supported by the National Natural Science Foundation of P.R. China (Grant No.12171064), by The team project of innovation leading talent in chongqing (No.CQYC20210309536) and by the Foundation of Chongqing Normal university (20XLB009); The second author's work was supported by the National Natural Science Foundation of P.R. China (Grant No.12001366), by the Science and Technology Research Program of Chongqing Municipal Education Commission(Grant No. KJZD-K202200506), by Chongqing Talent Program Lump-sum Project(No. cstc2022ycjh-bgzxm0040) and by the Foundation of Chongqing Normal University (22XLB005, ncamc2022-msxm02).

<sup>†</sup>Corresponding author. School of Mathematical Sciences, Chongqing Normal University, Chongqing 401331 P.R. China, Email: yisheng.song@cqnu.edu.cn

<sup>‡</sup>National Center for Applied Mathematics, Chongqing Normal University, Chongqing 401331 P.R. China, Email: jinjie.liu@cqnu.edu.cn

[1–6]. In the graph theory, the  $m$ th order copositive tensors may be directly applied to estimate the bounds on the independent number of  $m$ -uniform hypergraph [3, 7–9]. The concept of copositive tensors was introduced by Qi [10] in 2013, which is usually applied to a symmetric tensor or, more precisely, to its associated Homogeneous polynomial of degree  $m$ .

**Definition 1.1.** Let  $\mathcal{T} = (t_{i_1 i_2 \dots i_m})$  be an  $m$ th order  $n$  dimensional symmetric tensor.  $\mathcal{T}$  is called

- (i) **positive semi-definite** ([11]) if  $m$  is an even number and in the Euclidean space  $\mathbb{R}^n$ , its associated Homogeneous polynomial

$$\mathcal{T}x^m = \sum_{i_1, i_2, \dots, i_m=1}^n t_{i_1 i_2 \dots i_m} x_{i_1} x_{i_2} \cdots x_{i_m} \geq 0;$$

- (ii) **positive definite** ([11]) if  $m$  is an even number and  $\mathcal{T}x^m > 0$  for all  $x \in \mathbb{R}^n \setminus \{0\}$ ;

- (iii) **copositive** ([10]) if  $\mathcal{T}x^m \geq 0$  on the nonnegative orthant  $\mathbb{R}_+^n$ ;

- (iv) **strictly copositive** ([10]) if  $\mathcal{T}x^m > 0$  for all  $x \in \mathbb{R}_+^n \setminus \{0\}$ .

Clearly, the positive semi-definite tensors must be copositive, and a copositive tensor coincides with a copositive matrix if  $m = 2$ . The concept of copositive matrices was introduced by Motzkin [12] in 1952. Baston [13] gave an analytic way of judging copositivity of a  $n \times n$  matrix in 1969.

**Theorem 1.1.** (Baston [13]) Let  $M = (m_{ij})$  be a symmetric matrix with  $|m_{ij}| = m_{ii} = 1$  for all  $i, j \in \{1, 2, \dots, n\}$ . Then the matrix  $M$  is copositive if and only if there is no triple  $(r, s, t)$  such that

$$m_{rs} = m_{rt} = m_{st} = -1.$$

Simpson-Spector [14] and Hadeler [15] and Nadler [16] and Chang-Sederberg [17] and Andersson-Chang-Elfving [18] respectively showed the (strict) copositive conditions of  $2 \times 2$  and  $3 \times 3$  matrices using different methods of argumentation.

**Theorem 1.2.** Let  $M = (m_{ij})$  be a symmetric matrix. Then a  $2 \times 2$  matrix  $M$  is (strictly) copositive if and only if

$$m_{11} \geq 0 (> 0), m_{22} \geq 0 (> 0), m_{12} + \sqrt{m_{11}m_{22}} \geq 0 (> 0);$$

a  $3 \times 3$  matrix  $M$  is (strictly) copositive if and only if for all  $i \in \{1, 2, 3\}$ ,

$$\begin{aligned} m_{ii} &\geq 0 (> 0), \alpha = m_{12} + \sqrt{m_{11}m_{22}} \geq 0 (> 0), \\ \beta &= m_{13} + \sqrt{m_{11}m_{33}} \geq 0 (> 0), \gamma = m_{23} + \sqrt{m_{33}m_{22}} \geq 0 (> 0), \\ m_{12}\sqrt{m_{33}} + m_{13}\sqrt{m_{22}} + m_{23}\sqrt{m_{11}} + \sqrt{m_{11}m_{22}m_{33}} + \sqrt{2\alpha\beta\gamma} &\geq 0 (> 0). \end{aligned}$$

Schmidt-Heß [19] provided the nonnegative conditions of a cubic and univariate polynomial with real coefficients in non-negative real number  $\mathbb{R}_+$ , and Qi-Song-Zhang [20] recently gave the positivity of such a cubic polynomial, which actually gave a the (strict) copositivity of 3rd order 2-dimensional symmetric tensor (see Liu-Song [21] for more details also). Qi-Song-Zhang [20] also gave the nonnegativity and positivity of a quartic and univariate polynomial in  $\mathbb{R}$ , which means the positive (semi-)definiteness of 4th order 2-dimensional tensor. Ulrich-Watson [22] and Qi-Song-Zhang [20] presented the analytic conditions of the nonnegativity of a quartic and univariate polynomial in  $\mathbb{R}_+$ . This actually yielded the copositivity of 4th order 2-dimensional symmetric tensor [4].

**Theorem 1.3.** *A 3rd order 2-dimensional symmetric tensor  $\mathcal{T} = (t_{ijk})$  is (strictly) copositive if and only if  $t_{111} \geq 0$  ( $> 0$ ),  $t_{222} \geq 0$  ( $> 0$ ), either  $t_{112} \geq 0$ ,  $t_{122} \geq 0$  or*

$$4t_{111}t_{122}^3 + 4t_{112}^3t_{222} + t_{111}^2t_{222}^2 - 6t_{111}t_{112}t_{122}t_{222} - 3t_{112}^2t_{122}^2 \geq 0 \text{ (} > 0 \text{)}.$$

*A 4th order 2-dimensional symmetric tensor  $\mathcal{T} = (t_{ijkl})$  with  $t_{1111} > 0$  and  $t_{2222} > 0$  is copositive if and only if*

$$\begin{cases} \Delta \leq 0, t_{1222}\sqrt{t_{1111}} + t_{1112}\sqrt{t_{2222}} > 0; \\ t_{1222} \geq 0, t_{1112} \geq 0, 3t_{1122} + \sqrt{t_{1111}t_{2222}} \geq 0; \\ \Delta \geq 0, \\ |t_{1112}\sqrt{t_{2222}} - t_{1222}\sqrt{t_{1111}}| \leq \sqrt{6t_{1111}t_{1122}t_{2222} + 2t_{1111}t_{2222}\sqrt{t_{1111}t_{2222}}} \\ (i) -\sqrt{t_{1111}t_{2222}} \leq 3t_{1122} \leq 3\sqrt{t_{1111}t_{2222}}; \\ (ii) t_{1122} > \sqrt{t_{1111}t_{2222}} \text{ and} \\ t_{1112}\sqrt{t_{2222}} + t_{1222}\sqrt{t_{1111}} \geq -\sqrt{6t_{1111}t_{1122}t_{2222} - 2t_{1111}t_{2222}\sqrt{t_{1111}t_{2222}}}, \end{cases}$$

where  $\Delta = 4 \times 12^3(t_{1111}t_{2222} - 4t_{1112}t_{1222} + 3t_{1122}^2)^3 - 72^2 \times 6^2(t_{1111}t_{1122}t_{2222} + 2t_{1112}t_{1122}t_{1222} - t_{1122}^3 - t_{1112}^2t_{2222} - t_{1111}t_{1222}^2)^2$ .

For a special 4th order 3-dimensional tensor given by the particle physical model, Qi-Song-Zhang [23] presented a necessary and sufficient condition of copositivity, and Song-Li [4] provided an analytic condition of its copositivity. However, an analytic necessary and sufficient conditions has not been found for the copositivity of a general 3-dimensional higher order tensor ( $m > 2$ ). Even for a general 2-dimensional higher order tensor ( $m > 3$ ), people still has not found the analytic conditions of strict copositivity untill now.

For checking the copositivity of symmetric tensors, various numerical algorithms have been employed. For example, Chen-Huang-Qi [3, 24] gave the detection algorithms based on simplicial partition; Li-Zhang-Huang-Qi [25] used an SDP relaxation algorithm; Nie-Yang-Zhang [7] devised a complete semi-definite algorithms. Taking advantage of the properties of a tensor itself, the copositivity can be

described qualitatively. Song-Qi [26] showed a necessary and sufficient condition of copositivity by means of the principal sub-tensors; Song-Qi [27] applied the sign of its Pareto H-eigenvalue (Z-eigenvalue) to test the copositivity. Song-Qi [28] proved the equivalence of (strict) copositivity and (strict) semi-positivity of a symmetric tensor.

**Theorem 1.4.** (Song-Qi [28]) *Let  $\mathcal{T} = (t_{i_1 i_2 \dots i_m})$  be a symmetric tensor. Then  $\mathcal{T}$  is (strictly) copositive if and only if  $\mathcal{T}$  is (strictly) semi-positive, i.e., for each  $x = (x_1, x_2, \dots, x_n)^\top \in \mathbb{R}_+^n \setminus \{0\}$ , there exists  $k \in \{1, 2, \dots, n\}$  such that*

$$x_k > 0 \text{ and } (\mathcal{T}x^{m-1})_k = \sum_{i_2 \dots i_m=1}^n t_{k i_2 \dots i_m} x_{i_2} \dots x_{i_m} \geq 0 (> 0).$$

For more details of the copositivity of a higher order tensor and a matrix, see [8, 29–37]. Since there are strong connection between the semi-positivity of a tensor and the tensor complementarity problems [26, 38–40], so the copositivity of a symmetric tensor may be verified by solving the corresponding tensor complementarity problems. For more details about the tensor complementarity problems and its applications, see Refs. [41–59].

Motivation to checking the copositivity of a higher order tensor, we mainly discuss analytic necessary and sufficient conditions of copositivity of a class of 4th order 3-dimensional symmetric tensors in this paper. With the help of Theorem 1.4, we first promote Theorem 1.1 to ones of 4th order 3-dimensional symmetric tensors, which gives an analytic necessary and sufficient condition of strict copositivity of such a class of tensors (Theorem 3.4). Secondly, we present an analytic necessary and sufficient condition of copositivity of such a class of tensors (Theorem 3.8). Finally, applying Theorems 3.4 and 3.8, the analytic sufficient conditions (Theorems 3.9, 3.10 and 3.11) are successfully proved for (strict) copositivity of a general 4th order 3-dimensional symmetric tensor also.

## 2 Copositivity of 4th order 2-dimensional symmetric tensors

Let  $T = (t_{i_1 \dots i_m})(i_j = 1, 2, \dots, n, j = 1, 2, \dots, m)$  be a  $m$ th-order  $n$ -dimensional symmetric tensor. Then for  $x = (x_1, x_2, \dots, x_n)^\top \in \mathbb{R}^n$ , we write

$$Tx^m = x^\top (Tx^{m-1}) = \sum_{i_1 \dots i_m=1}^n t_{i_1 \dots i_m} x_{i_1} \dots x_{i_m},$$

and  $Tx^{m-1} = (y_1, y_2, \dots, y_n)^\top$  is a vector with its components

$$y_k = (Tx^{m-1})_k = \sum_{i_2 \dots i_m=1}^n t_{ki_2 \dots i_m} x_{i_2} \dots x_{i_m}, \quad k = 1, 2, \dots, n.$$

Let  $f(x_1, x_2)$  be a quartic homogeneous real polynomial about two variables  $x_1, x_2$ ,

$$f(x_1, x_2) = x_1^4 + 4ax_1^3x_2 + 6bx_1^2x_2^2 + 4cx_1x_2^3 + x_2^4. \quad (1)$$

Then it gives a 4th-order 2-dimensional symmetric tensor  $\mathcal{T} = (t_{ijkl})$  with its entries,

$$t_{1111} = 1, t_{1112} = a, t_{1122} = b, t_{1222} = c, t_{2222} = 1. \quad (2)$$

By Theorem 1.3, the following lemma can be obtained easily.

**Lemma 2.1.** *Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 2-dimensional symmetric tensor given by (2). Then  $\mathcal{T}$  is copositive, i.e.,  $f(x_1, x_2) \geq 0$  for all  $x = (x_1, x_2)^\top \geq 0$  if and only if*

- (1)  $\Delta' \leq 0$  and  $a + c > 0$ ;
- (2)  $a \geq 0, c \geq 0$  and  $1 + 3b \geq 0$ ;
- (3)  $\Delta' \geq 0, |a - c| \leq \sqrt{6b + 2}$  and
  - (i)  $-1 \leq 3b \leq 3$ ,
  - (ii)  $b > 1$  and  $a + c \geq -\sqrt{6b - 2}$ ,

where  $\Delta = 4 \times 12^3 \Delta', \Delta' = (1 - 4ac + 3b^2)^3 - 27(b + 2abc - b^3 - c^2 - a^2)^2$ .

**Lemma 2.2.** *Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 2-dimensional symmetric tensor with its entries  $|t_{ijkl}| = 1$  and  $t_{1111} = t_{2222} = 1$ . Then  $\mathcal{T}$  is copositive if and only if either*

$$b = t_{1122} = 1 \text{ or } a = t_{1112} = t_{1222} = c = 1.$$

*Proof.* It follows from Lemma 2.1 that  $\mathcal{T}$  is copositive if and only if

- (1)  $\Delta' \leq 0$  and  $a + c > 0$ ;
- (2)  $a \geq 0, c \geq 0$  and  $1 + 3b \geq 0$ ;
- (3)  $\Delta' \geq 0, |a - c| \leq \sqrt{6b + 2}$  and  $-1 \leq 3b \leq 3$ .

Since  $|t_{ijkl}| = 1$ , then the conditions (1)-(3) mean

- (1)  $\Delta' \leq 0$  and  $a = c = 1 \Leftrightarrow a = c = 1$  and either

$$b = 1, \Delta' = (1 - 4 + 3)^3 - 27(1 + 2 - 1^3 - 1 - 1)^2 = 0,$$

or

$$b = -1, \Delta' = (1 - 4 + 3)^3 - 27(-1 - 2 + 1 - 1 - 1)^2 < 0;$$

(2)  $a = c = 1$  and  $b = 1$ ;

(3)  $\Delta' \geq 0$ ,  $|a - c| \leq \sqrt{6b + 2}$  and  $b = 1 \Leftrightarrow b = 1$  and either  $ac = 1$ ,  
 $\Delta' = (1 - 4 + 3)^3 - 27(1 + 2 - 1 - 1 - 1)^2 = 0$ ,  $|a - c| = 0 < \sqrt{6b + 2} = \sqrt{8}$ ;  
or  $ac = -1$ ,

$\Delta' = (1 + 4 + 3)^3 - 27(1 - 2 - 1 - 1 - 1)^2 > 0$ ,  $|a - c| = 2 < \sqrt{6b + 2} = \sqrt{8}$ .

So the conditions (1)-(3) are equivalent to

$$b = 1 \text{ or } a = c = 1.$$

This completes the proof.  $\square$

**Corollary 2.1.** *Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 2-dimensional symmetric tensor with its entires  $t_{1111} \geq 1$  and  $t_{2222} \geq 1$ . Then*

(1)  $\mathcal{T}$  is strictly copositive if

$$t_{1112} \geq 1, t_{1222} \geq 1, t_{1122} \geq -1;$$

(2)  $\mathcal{T}$  is copositive if

$$t_{1112} \geq -1, t_{1222} \geq -1, t_{1122} \geq 1.$$

*Proof.* Let  $x = (x_1, x_2)^\top \geq 0$ . Then

$$\mathcal{T}x^4 = t_{1111}x_1^4 + 4t_{1112}x_1^3x_2 + 6t_{1122}x_1^2x_2^2 + 4t_{1222}x_1x_2^3 + t_{2222}x_2^4.$$

(1) It is obvious that

$$\begin{aligned} \mathcal{T}x^4 &\geq x_1^4 + 4x_1^3x_2 - 6x_1^2x_2^2 + 4x_1x_2^3 + x_2^4 \\ &= (x_1^2 + x_2^2)^2 + 4x_1x_2(x_1^2 - 2x_1x_2 + x_2^2) \\ &= (x_1^2 + x_2^2)^2 + 4x_1x_2(x_1 - x_2)^2. \end{aligned}$$

So  $\mathcal{T}x^4 > 0$  for  $x \geq 0$  and  $x \neq 0$ . Suppose not, then there exists  $x = (x_1, x_2)^\top \neq 0$  such that  $\mathcal{T}x^4 = 0$ , and hence,

$$0 = \mathcal{T}x^4 \geq (x_1^2 + x_2^2)^2 + 4x_1x_2(x_1 - x_2)^2 \geq 0.$$

That is,

$$(x_1^2 + x_2^2)^2 + 4x_1x_2(x_1 - x_2)^2 = 0,$$

which means  $x_1^2 + x_2^2 = 0$ , i.e.,  $x_1 = x_2 = 0$ , a contradiction. Therefore,  $\mathcal{T}$  is strictly copositive.

(2) For any  $x \geq 0$ , it follows from Lemma 2.2 that

$$\mathcal{T}x^4 \geq x_1^4 - 4x_1^3x_2 + 6x_1^2x_2^2 - 4x_1x_2^3 + x_2^4 \geq 0.$$

So,  $\mathcal{T}$  is copositive. This completes the proof.  $\square$

For a 4th-order 2-dimensional symmetric tensor  $\mathcal{A} = (a_{ijkl})$  with its entires  $a_{1111} > 0$  and  $a_{2222} > 0$ , let  $t_{1111} = t_{2222} = 1$  and

$$t_{1112} = a_{1112}a_{1111}^{-\frac{3}{4}}a_{2222}^{-\frac{1}{4}}, t_{1122} = a_{1122}a_{1111}^{-\frac{1}{2}}a_{2222}^{-\frac{1}{2}}, t_{1222} = a_{1222}a_{1111}^{-\frac{1}{4}}a_{2222}^{-\frac{3}{4}}.$$

For  $y = (y_1, y_2)^\top$  and  $x = (x_1, x_2)^\top = (a_{1111}^{\frac{1}{4}}y_1, a_{2222}^{\frac{1}{4}}y_2)^\top$ , then

$$\begin{aligned}\mathcal{A}y^4 &= a_{1111}y_1^4 + 4a_{1112}y_1^3y_2 + 6a_{1122}y_1^2y_2^2 + 4a_{1222}y_1y_2^3 + a_{2222}y_2^4 \\ &= x_1^4 + 4t_{1112}x_1^3x_2 + 6t_{1122}x_1^2x_2^2 + 4t_{1222}x_1x_2^3 + x_2^4 \\ &= \mathcal{T}x^4.\end{aligned}$$

Obviously, the copositivity of  $\mathcal{A} = (a_{ijkl})$  coincides with one of  $\mathcal{T} = (t_{ijkl})$ , and hence, we can establish an analytically sufficient condition of the (strict) copositivity of a gerenal 4th-order 2-dimensional symmetric tensor  $\mathcal{A} = (a_{ijkl})$ , that can be very easily parsed and validated.

**Corollary 2.2.** *Let  $\mathcal{A} = (a_{ijkl})$  be a 4th-order 2-dimensional symmetric tensor with its entires  $a_{1111} > 0$  and  $a_{2222} > 0$ . Then*

(1)  $\mathcal{A}$  is strictly copositive if

$$a_{1112} \geq a_{1111}^{\frac{3}{4}}a_{2222}^{\frac{1}{4}}, a_{1122} \geq -\sqrt{a_{1111}a_{2222}}, a_{1222} \geq a_{1111}^{\frac{1}{4}}a_{2222}^{\frac{3}{4}};$$

(2)  $\mathcal{A}$  is copositive if

$$a_{1112} \geq -a_{1111}^{\frac{3}{4}}a_{2222}^{\frac{1}{4}}, a_{1122} \geq \sqrt{a_{1111}a_{2222}}, a_{1222} \geq -a_{1111}^{\frac{1}{4}}a_{2222}^{\frac{3}{4}}.$$

### 3 Copositivity of 4th order 3-dimensional symmetric tensors

#### 3.1 Analytical expressions of strict copositivity

**Theorem 3.1.** Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 3-dimensional symmetric tensor with  $|t_{ijkl}| = t_{iiii} = t_{iiij} = 1$  for all  $i, j, k, l \in \{1, 2, 3\}$ . If there is at most one  $-1$  in  $\{t_{1123}, t_{1223}, t_{1233}\}$ , then  $\mathcal{T}$  is strictly copositive.

*Proof.* Without loss the generality, let  $t_{1123} = -1, t_{1223} = t_{1233} = 1$ . Then

$$\begin{aligned}\mathcal{T}x^4 &= x_1^4 + x_2^4 + x_3^4 + 6t_{1122}x_1^2x_2^2 + 6t_{1133}x_1^2x_3^2 + 6t_{2233}x_2^2x_3^2 \\ &\quad + 4x_1^3x_2 + 4x_1^3x_3 + 4x_1x_2^3 + 4x_1x_3^3 + 4x_2^3x_3 + 4x_2x_3^3 \\ &\quad - 12x_1^2x_2x_3 + 12x_1x_2^2x_3 + 12x_1x_2x_3^2,\end{aligned}$$

and so,

$$\mathcal{T}x^3 = \frac{1}{4}\nabla\mathcal{T}x^4 = \begin{pmatrix} \sum_{j,k,l=1}^3 t_{1jkl}x_jx_kx_l \\ \sum_{j,k,l=1}^3 t_{2jkl}x_jx_kx_l \\ \sum_{j,k,l=1}^3 t_{3jkl}x_jx_kx_l \end{pmatrix}$$

It follows from Theorem 1.4 that we need only show that  $\mathcal{T} = (t_{ijkl})$  is strictly semi-positive, i.e., for  $x = (x_1, x_2, x_3)^\top \geq 0$ , there exists  $k \in \{1, 2, 3\}$  such that

$$x_k > 0 \text{ and } (\mathcal{T}x^3)_k > 0.$$

For each  $i \in \{1, 2, 3\}$ , we have

$$\begin{aligned} (\mathcal{T}x^3)_1 &= \sum_{j,k,l=1}^3 t_{1jkl}x_jx_kx_l = x_1^3 + x_2^3 + x_3^3 + 3t_{1122}x_1x_2^2 + 3t_{1133}x_1x_3^2 \\ &\quad + 3x_1^2x_2 + 3x_1^2x_3 + 3x_2^2x_3 + 3x_2x_3^2 - 6x_1x_2x_3 \\ &\geq x_1^3 + x_2^3 + x_3^3 - 3x_1x_2^2 - 3x_1x_3^2 + 3x_1^2x_2 \\ &\quad + 3x_1^2x_3 + 3x_2^2x_3 + 3x_2x_3^2 - 6x_1x_2x_3 \\ &= (x_2 + x_3 - x_1)^3 + 2x_1^3 = x_1^3 + (x_2 + x_3 - x_1 + x_1) \\ &\quad \times ((x_2 + x_3 - x_1)^2 - x_1(x_2 + x_3 - x_1) + x_1^2) \\ &= x_1^3 + (x_2 + x_3)((x_2 + x_3 - x_1)^2 - x_1(x_2 + x_3 - x_1) + x_1^2); \\ (\mathcal{T}x^3)_2 &= \sum_{j,k,l=1}^3 t_{2jkl}x_jx_kx_l = x_1^3 + x_2^3 + x_3^3 + 3t_{1122}x_1^2x_2 + 3t_{2233}x_2x_3^2 \\ &\quad + 3x_1x_2^2 + 3x_2^2x_3 - 3x_1^2x_3 + 3x_1x_3^2 + 6x_1x_2x_3 \\ &\geq x_1^3 + x_2^3 + x_3^3 - 3x_1^2x_2 - 3x_2x_3^2 + 3x_1x_2^2 \\ &\quad + 3x_2^2x_3 - 3x_1^2x_3 + 3x_1x_3^2 + 6x_1x_2x_3 \\ &= (x_1 + x_3 - x_2)^3 + 2x_2^3 + 12x_1x_2x_3 - 6x_1^2x_3 \\ &= ((x_1 + x_3 - x_2)^3 + 2x_2^3) + 6x_1x_3(2x_2 - x_1); \end{aligned}$$



$$\begin{aligned}
(\mathcal{T}x^3)_3 &= \sum_{j,k,l=1}^3 t_{3jkl}x_jx_kx_l = x_1^3 + x_2^3 + x_3^3 + 3t_{1133}x_1^2x_3 + 3t_{2233}x_2^2x_3 \\
&\quad + 3x_1x_3^2 + 3x_2x_3^2 - 3x_1^2x_2 + 3x_1x_2^2 + 6x_1x_2x_3 \\
&\geq x_1^3 + x_2^3 + x_3^3 - 3x_1^2x_3 - 3x_2^2x_3 + 3x_1x_3^2 \\
&\quad + 3x_2x_3^2 - 3x_1^2x_2 + 3x_1x_2^2 + 6x_1x_2x_3 \\
&= (x_1 + x_2 - x_3)^3 + 2x_3^3 + 12x_1x_2x_3 - 6x_1^2x_2 \\
&= ((x_1 + x_2 - x_3)^3 + 2x_3^3) + 6x_1x_2(2x_3 - x_1).
\end{aligned}$$

So, it follows that

- $x_1 > 0$ ,  $(\mathcal{T}x^3)_1 > 0$ , which is done; otherwise,
- $x_1 = 0, x_2 > 0$ ,  $(\mathcal{T}x^3)_2 > 0$ ;
- $x_1 = 0, x_3 > 0$ ,  $(\mathcal{T}x^3)_3 > 0$ ,

and hence,  $\mathcal{T}$  is strictly copositive.  $\square$

We definite

$$\mathcal{T}' = (t'_{ijkl}) \leq \mathcal{T} = (t_{ijkl}) \Leftrightarrow t'_{ijkl} \leq t_{ijkl}, \text{ for all } i, j, k, l.$$

Then for all  $x \in \mathbb{R}_+^n$ , we have

$$\mathcal{T}'x^4 = \sum_{i,j,k,l=1}^n t'_{ijkl}x_ix_jx_kx_l \leq \sum_{i,j,k,l=1}^n t_{ijkl}x_ix_jx_kx_l = \mathcal{T}x^4.$$

So, the (strict) copositivity of a tensor  $\mathcal{T}'$  implies one of  $\mathcal{T}$ . Therefore, from Theorem 3.1, the following conclusions are obvious.

**Corollary 3.2.** Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 3-dimensional symmetric tensor. If  $t_{iiii} \geq 1, t_{iiij} \geq 1, t_{iijj} \geq -1$  for all  $i, j \in \{1, 2, 3\}, i \neq j$  and one of the following conditions,

- (1)  $t_{1123} \geq -1, t_{1223} \geq 1, t_{1233} \geq 1$ ;
- (2)  $t_{1123} \geq 1, t_{1223} \geq -1, t_{1233} \geq 1$ ;
- (3)  $t_{1123} \geq 1, t_{1223} \geq 1, t_{1233} \geq -1$ ,

then  $\mathcal{T}$  is strictly copositive.

**Theorem 3.3.** Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 3-dimensional symmetric tensor with  $|t_{ijkl}| = t_{iiii} = t_{iiij} = 1$  for all  $i, j, k, l \in \{1, 2, 3\}$ . Then  $\mathcal{T}$  is strictly copositive if and only if

- (1)  $t_{1122} = t_{1133} = t_{2233} = 1$  if  $t_{1123} = t_{1223} = t_{1233} = -1$ ;
- (2) there is at least one 1 in  $\{t_{1122}, t_{1133}, t_{2233}\}$  if two of  $\{t_{1123}, t_{1223}, t_{1233}\}$  are only  $-1$ .

*Proof.* Necessity. If  $\mathcal{T}$  is strictly copositive, but the conditions don't hold. Then for  $x = (1, 1, 1)^\top$ , it follows that (1) one of  $\{t_{1122}, t_{1133}, t_{2233}\}$  is  $-1$ ,

$$\begin{aligned}\mathcal{T}x^4 &= x_1^4 + x_2^4 + x_3^4 + 6t_{1122}x_1^2x_2^2 + 6t_{1133}x_1^2x_3^2 + 6t_{2233}x_2^2x_3^2 \\ &\quad + 4x_1^3x_2 + 4x_1^3x_3 + 4x_1x_2^3 + 4x_1x_3^3 + 4x_2^3x_3 + 4x_2x_3^3 \\ &\quad + 12t_{1123}x_1^2x_2x_3 + 12t_{1223}x_1x_2^2x_3 + 12t_{1233}x_1x_2x_3^2 \\ &= 27 + 12 - 6 - 36 = -3 < 0;\end{aligned}$$

$$(2) \ t_{1122} = t_{1133} = t_{2233} = -1,$$

$$\mathcal{T}x^4 = \sum_{i,j,k,l=1}^3 t_{ijkl}x_ix_jx_kx_l = 27 - 18 - 24 + 12 = -3 < 0.$$

So,  $\mathcal{T}$  is not strictly copositive.

Sufficiency. (1) Since  $t_{1122} = t_{1133} = t_{2233} = 1$  and  $t_{1123} = t_{1223} = t_{1233} = -1$ , then

$$\begin{aligned}\mathcal{T}x^4 &= x_1^4 + x_2^4 + x_3^4 + 6x_1^2x_2^2 + 6x_1^2x_3^2 + 6x_2^2x_3^2 \\ &\quad + 4x_1^3x_2 + 4x_1^3x_3 + 4x_1x_2^3 + 4x_1x_3^3 + 4x_2^3x_3 + 4x_2x_3^3 \\ &\quad - 12x_1^2x_2x_3 - 12x_1x_2^2x_3 - 12x_1x_2x_3^2,\end{aligned}$$

and so,

$$\begin{aligned}(\mathcal{T}x^3)_1 &= \sum_{j,k,l=1}^3 t_{1jkl}x_jx_kx_l \\ &= x_1^3 + x_2^3 + x_3^3 + 3x_1x_2^2 + 3x_1x_3^2 + 3x_1^2x_2 + 3x_1^2x_3 \\ &\quad - 3x_2^2x_3 - 3x_2x_3^2 - 6x_1x_2x_3 \\ &= (x_2 + x_3 - x_1)^3 + 2x_1^3 + 6x_1x_2^2 + 6x_1x_3^2 - 6x_2^2x_3 - 6x_2x_3^2 \\ &= (x_2 + x_3 - x_1)^3 + 2x_1^3 + 6x_2^2(x_1 - x_3) + 6x_3^2(x_1 - x_2);\end{aligned}$$

$$\begin{aligned}(\mathcal{T}x^3)_2 &= \sum_{j,k,l=1}^3 t_{2jkl}x_jx_kx_l \\ &= x_1^3 + x_2^3 + x_3^3 + 3x_1^2x_2 + 3x_2x_3^2 + 3x_1x_2^2 + 3x_2^2x_3 \\ &\quad - 3x_1^2x_3 - 6x_1x_2x_3 - 3x_1x_3^2 \\ &= (x_1 + x_3 - x_2)^3 + 2x_2^3 + 6x_1^2x_2 + 6x_2x_3^2 - 6x_1^2x_3 - 6x_1x_3^2 \\ &= (x_1 + x_3 - x_2)^3 + 2x_2^3 + 6x_1^2(x_2 - x_3) + 6x_3^2(x_2 - x_1);\end{aligned}$$

$$\begin{aligned}
(\mathcal{T}x^3)_3 &= \sum_{j,k,l=1}^3 t_{3jkl}x_jx_kx_l \\
&= x_1^3 + x_2^3 + x_3^3 + 3x_1^2x_3 + 3x_2^2x_3 + 3x_1x_3^2 + 3x_2x_3^2 \\
&\quad - 3x_1^2x_2 - 3x_1x_2^2 - 6x_1x_2x_3 \\
&= (x_1 + x_2 - x_3)^3 + 2x_3^3 + 6x_1^2x_3 + 6x_2^2x_3 - 6x_1^2x_2 - 6x_1x_2^2 \\
&= (x_1 + x_2 - x_3)^3 + 2x_3^3 + 6x_1^2(x_3 - x_2) + 6x_2^2(x_3 - x_1).
\end{aligned}$$

So, it follows that

- $x_1 \geq \max\{x_2, x_3\}$ ,  $(\mathcal{T}x^3)_1 > 0$ , which is done; otherwise,
- $x_2 \geq \max\{x_1, x_3\}$ ,  $(\mathcal{T}x^3)_2 > 0$ ;
- $x_3 \geq \max\{x_1, x_2\}$ ,  $(\mathcal{T}x^3)_3 > 0$ ;

and hence,  $\mathcal{T}$  is strictly copositive.

(2) We might take  $t_{1122} = t_{1233} = 1$  and  $t_{1133} = t_{2233} = t_{1123} = t_{1223} = -1$ , then

$$\begin{aligned}
\mathcal{T}x^4 &= x_1^4 + x_2^4 + x_3^4 + 6x_1^2x_2^2 - 6x_1^2x_3^2 - 6x_2^2x_3^2 \\
&\quad + 4x_1^3x_2 + 4x_1^3x_3 + 4x_1x_2^3 + 4x_1x_3^3 + 4x_2^3x_3 + 4x_2x_3^3 \\
&\quad - 12x_1^2x_2x_3 - 12x_1x_2^2x_3 + 12x_1x_2x_3^2,
\end{aligned}$$

and so,

$$\begin{aligned}
(\mathcal{T}x^3)_1 &= \sum_{j,k,l=1}^3 t_{1jkl}x_jx_kx_l \\
&= x_1^3 + x_2^3 + x_3^3 + 3x_1x_2^2 - 3x_1x_3^2 + 3x_1^2x_2 + 3x_1^2x_3 \\
&\quad - 3x_2^2x_3 + 3x_2x_3^2 - 6x_1x_2x_3 \\
&= (x_2 + x_3 - x_1)^3 + 2x_1^3 + 6x_1x_2^2 - 6x_2^2x_3 \\
&= (x_2 + x_3 - x_1)^3 + 2x_1^3 + 6x_2^2(x_1 - x_3);
\end{aligned}$$

$$\begin{aligned}
(\mathcal{T}x^3)_2 &= \sum_{j,k,l=1}^3 t_{2jkl}x_jx_kx_l \\
&= x_1^3 + x_2^3 + x_3^3 + 3x_1^2x_2 - 3x_2^2x_3 + 3x_1x_2^2 + 3x_2^2x_3 \\
&\quad - 3x_1^2x_3 - 6x_1x_2x_3 + 3x_1x_3^2 \\
&= (x_1 + x_3 - x_2)^3 + 2x_2^3 + 6x_1^2x_2 - 6x_1^2x_3 \\
&= (x_1 + x_3 - x_2)^3 + 2x_2^3 + 6x_1^2(x_2 - x_3);
\end{aligned}$$

$$\begin{aligned}
(\mathcal{T}x^3)_3 &= \sum_{j,k,l=1}^3 t_{3jkl}x_jx_kx_l \\
&= x_1^3 + x_2^3 + x_3^3 - 3x_1^2x_3 - 3x_2^2x_3 + 3x_1x_3^2 + 3x_2x_3^2 \\
&\quad - 3x_1^2x_2 - 3x_1x_2^2 + 6x_1x_2x_3 \\
&= (x_1 + x_2 - x_3)^3 + 2x_3^3 + 12x_1x_2x_3 - 6x_1^2x_2 - 6x_1x_2^2 \\
&= (x_1 + x_2 - x_3)^3 + 2x_3^3 + 6x_1x_2((x_3 - x_2) + (x_3 - x_1)).
\end{aligned}$$

So, it follows that

- $x_1 \geq x_3$  and  $x_1 > 0$ ,  $(\mathcal{T}x^3)_1 > 0$ , which is done; otherwise,
- $x_2 \geq x_3$  and  $x_2 > 0$ ,  $(\mathcal{T}x^3)_2 > 0$ ;
- $x_3 \geq \max\{x_1, x_2\}$ ,  $(\mathcal{T}x^3)_3 > 0$ ;

and hence,  $\mathcal{T}$  is strictly copositive.  $\square$

Combing the conclusions of Theorems 3.1 and 3.3, the main result is bulit in this subsection.

**Theorem 3.4.** Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 3-dimensional symmetric tensor with  $|t_{ijkl}| = t_{iiii} = t_{iiij} = 1$  for all  $i, j, k, l \in \{1, 2, 3\}$ . Then  $\mathcal{T}$  is strictly copositive if and only if

- (1) there is at most one  $-1$  in  $\{t_{1123}, t_{1223}, t_{1233}\}$ ;
- (2) two of  $\{t_{1123}, t_{1223}, t_{1233}\}$  are only  $-1$  and there is at least one  $1$  in  $\{t_{1122}, t_{1133}, t_{2233}\}$ ;
- (3)  $t_{1123} = t_{1223} = t_{1233} = -1$  and  $t_{1122} = t_{1133} = t_{2233} = 1$ .

### 3.2 Analytical expressions of copositivity

**Theorem 3.5.** Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 3-dimensional symmetric tensor with  $|t_{ijkl}| = t_{iiii} = t_{iiij} = 1$  for all  $i, j, k, l \in \{1, 2, 3\}$ . If there is at most one  $-1$  in  $\{t_{1123}, t_{1223}, t_{1233}\}$ , then  $\mathcal{T}$  is copositive.

*Proof.* Without loss the generality, let  $t_{1123} = -1, t_{1223} = t_{1233} = 1$ . Then

$$\begin{aligned}
\mathcal{T}x^4 &= x_1^4 + x_2^4 + x_3^4 + 6x_1^2x_2^2 + 6x_1^2x_3^2 + 6x_2^2x_3^2 + 4t_{1112}x_1^3x_2 + 4t_{1113}x_1^3x_3 \\
&\quad + 4t_{1222}x_1x_2^3 + 4t_{1333}x_1x_3^3 + 4t_{2223}x_2^3x_3 + 4t_{2333}x_2x_3^3 \\
&\quad - 12x_1^2x_2x_3 + 12x_1x_2^2x_3 + 12x_1x_2x_3^2,
\end{aligned}$$

and so,

$$\begin{aligned}
\mathcal{T}x^4 &\geq x_1^4 + x_2^4 + x_3^4 + 6x_1^2x_2^2 + 6x_1^2x_3^2 + 6x_2^2x_3^2 \\
&\quad - 4x_1^3x_2 - 4x_1^3x_3 - 4x_1x_2^3 - 4x_1x_3^3 - 4x_2^3x_3 - 4x_2x_3^3 \\
&\quad - 12x_1^2x_2x_3 + 12x_1x_2^2x_3 + 12x_1x_2x_3^2 \\
&= (x_1 + x_2 - x_3)^4 + 8(3x_1x_2^2x_3 - x_1^3x_2 - x_1x_3^2).
\end{aligned}$$

Let  $f(x_1, x_2, x_3) = (x_1 + x_2 - x_3)^4 + 8(3x_1x_2^2x_3 - x_1^3x_2 - x_1x_2^3)$ . Then the stationary points of the function  $f(x_1, x_2, x_3)$  are the solution to this system of equations,

$$\begin{cases} f'_{x_1}(x_1, x_2, x_3) = 4(x_1 + x_2 - x_3)^3 + 8(3x_2^2x_3 - 3x_1^2x_2 - x_2^3) = 0, \\ f'_{x_2}(x_1, x_2, x_3) = 4(x_1 + x_2 - x_3)^3 + 8(6x_1x_2x_3 - x_1^3 - 3x_1x_2^2) = 0, \\ f'_{x_3}(x_1, x_2, x_3) = -4(x_1 + x_2 - x_3)^3 + 24x_1x_2^2 = 0, \end{cases}$$

i.e.,

$$\begin{cases} 3x_1x_2^2 + 3x_2^2x_3 - 3x_1^2x_2 - x_2^3 = 0, \\ 3x_1x_2^2 + 6x_1x_2x_3 - x_1^3 - 3x_1x_2^2 = 0, \\ (x_1 + x_2 - x_3)^3 = 6x_1x_2^2, \end{cases}$$

and hence,

$$\begin{cases} x_2 = 0 \text{ or } 3x_1x_2 + 3x_2x_3 - 3x_1^2 - x_2^2 = 0, \\ x_1 = 0 \text{ or } 6x_2x_3 = x_1^2, \\ (x_1 + x_2 - x_3)^3 = 6x_1x_2^2. \end{cases}$$

So the stationary points are  $y = (0, 0, 0)$ , and hence,  $f(0, 0, 0) = 0$ . Since the boundary of the non-negative orthant  $\{(x_1, x_2, x_3); x_1 \geq 0, x_2 \geq 0, x_3 \geq 0\}$  are three coordinate planes, then at these boundary points, the function  $f(x_1, x_2, x_3)$  give three 4th-order 2-dimensional symmetric tensors. It follows from Lemma 2.2 that

$$f(0, x_2, x_3) \geq 0, f(x_1, 0, x_3) \geq 0, f(x_1, x_2, 0) \geq 0.$$

Thus  $\mathcal{T}x^4 \geq f(x_1, x_2, x_3) \geq 0$  for any  $x = (x_1, x_2, x_3)^\top \geq 0$ . That is,  $\mathcal{T}$  is copositive.  $\square$

From the proof of Theorem 3.5, the following conclusion is easily obtained.

**Corollary 3.6.** Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 3-dimensional symmetric tensor. If for all  $i, j \in \{1, 2, 3\}$  and  $i \neq j$ ,

$$t_{iiii} \geq 1, t_{iijj} \geq 1, t_{iiij} \geq -1$$

and one of the following three conditions holds,

- (a)  $t_{1123} \geq -1, t_{1223} \geq 1, t_{1233} \geq 1$ ;
- (b)  $t_{1123} \geq 1, t_{1223} \geq -1, t_{1233} \geq 1$ ;
- (c)  $t_{1123} \geq 1, t_{1223} \geq 1, t_{1233} \geq -1$ ,

then  $\mathcal{T}$  is copositive.

**Theorem 3.7.** Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 3-dimensional symmetric tensor with  $|t_{ijkl}| = t_{iiii} = t_{iijj} = 1$  for all  $i, j, k, l \in \{1, 2, 3\}$ . Then  $\mathcal{A}$  is copositive if and only if

- (1) there is at least two 1 in  $\{t_{iiij}; i, j = 1, 2, 3, i \neq j\}$  if there is only two  $-1$  in  $\{t_{1123}, t_{1223}, t_{1233}\}$ ;
- (2) there is at least five 1 in  $\{t_{iiij}; i, j = 1, 2, 3, i \neq j\}$  if  $t_{1123} = t_{1223} = t_{1233} = -1$ .

*Proof.* Necessity. If  $\mathcal{T}$  is copositive, but the conditions don't hold, then for  $x = (1, 1, 1)^\top$ ,

- (1) there are five  $-1$  in  $\{t_{iiij}; i, j = 1, 2, 3, i \neq j\}$ ,

$$\begin{aligned}\mathcal{T}x^4 &= x_1^4 + x_2^4 + x_3^4 + 6x_1^2x_2^2 + 6x_1^2x_3^2 + 6x_2^2x_3^2 \\ &\quad + 4t_{1112}x_1^3x_2 + 4t_{1113}x_1^3x_3 + 4t_{1222}x_1x_2^3 + 4t_{1333}x_1x_3^3 \\ &\quad + 4t_{2223}x_2^3x_3 + 4t_{2333}x_2x_3^3 \\ &\quad + 12t_{1123}x_1^2x_2x_3 + 12t_{1223}x_1x_2^2x_3 + 12t_{1233}x_1x_2x_3^2 \\ &= 21 + 4 - 20 - 24 + 12 = -7 < 0;\end{aligned}$$

- (2) there are two  $-1$  in  $\{t_{iiij}; i, j = 1, 2, 3, i \neq j\}$ ,

$$\mathcal{T}x^4 = \sum_{i,j,k,l=1}^3 t_{ijkl}x_i x_j x_k x_l = 21 + 16 - 8 - 36 = -7 < 0.$$

So,  $\mathcal{T}$  is not copositive, which is a contradiction, and hence, the conditions hold.

Sufficiency.  $\mathcal{T}x^4$  may be rewritten as follows,

$$\begin{aligned}\mathcal{T}x^4 &= (x_1 + x_2 - x_3)^4 + 4(t_{1112} - 1)x_1^3x_2 + 4(t_{1222} - 1)x_1x_2^3 \\ &\quad + 4(t_{1333} + 1)x_1x_3^3 + 4(t_{1113} + 1)x_1^3x_3 + 4(t_{2223} + 1)x_2^3x_3 \\ &\quad + 4(t_{2333} + 1)x_2x_3^3 + 12(t_{1123} + 1)x_1^2x_2x_3 \\ &\quad + 12(t_{1223} + 1)x_1x_2^2x_3 + 12(t_{1233} - 1)x_1x_2x_3^2;\end{aligned}$$

$$\begin{aligned}\mathcal{T}x^4 &= (x_1 - x_2 + x_3)^4 + 4(t_{1112} + 1)x_1^3x_2 + 4(t_{1222} + 1)x_1x_2^3 \\ &\quad + 4(t_{1333} - 1)x_1x_3^3 + 4(t_{1113} - 1)x_1^3x_3 + 4(t_{2223} + 1)x_2^3x_3 \\ &\quad + 4(t_{2333} + 1)x_2x_3^3 + 12(t_{1123} + 1)x_1^2x_2x_3 \\ &\quad + 12(t_{1223} - 1)x_1x_2^2x_3 + 12(t_{1233} + 1)x_1x_2x_3^2;\end{aligned}$$

$$\begin{aligned}\mathcal{T}x^4 &= (x_1 - x_2 - x_3)^4 + 4(t_{1112} + 1)x_1^3x_2 + 4(t_{1222} + 1)x_1x_2^3 \\ &\quad + 4(t_{1333} + 1)x_1x_3^3 + 4(t_{1113} + 1)x_1^3x_3 + 4(t_{2223} - 1)x_2^3x_3 \\ &\quad + 4(t_{2333} - 1)x_2x_3^3 + 12(t_{1123} - 1)x_1^2x_2x_3 \\ &\quad + 12(t_{1223} + 1)x_1x_2^2x_3 + 12(t_{1233} + 1)x_1x_2x_3^2.\end{aligned}$$

Clearly, for the boundary points of the non-negative orthant,  $x = (0, x_2, x_3), (x_1, 0, x_3), (x_1, x_2, 0)$ , it follows from Lemma 2.2 that  $\mathcal{T}x^4 \geq 0$ .

(1) Without loss the generality, let  $t_{1112} = t_{2223} = t_{1233} = 1$  and  $t_{1113} = t_{1333} = t_{2333} = t_{1222} = t_{1123} = t_{1223} = -1$ . Then

$$\mathcal{T}x^4 = (x_1 + x_2 - x_3)^4 + 8x_2^3(x_3 - x_1)$$

Let  $f(x_1, x_2, x_3) = \mathcal{T}x^4$ . Then the stationary points of the function  $f(x_1, x_2, x_3)$  are the solution to this system of equations,

$$\begin{cases} f'_{x_1}(x_1, x_2, x_3) = 4(x_1 + x_2 - x_3)^3 - 8x_2^3 = 0, \\ f'_{x_2}(x_1, x_2, x_3) = 4(x_1 + x_2 - x_3)^3 + 24x_2^2(x_3 - x_1) = 0, \\ f'_{x_3}(x_1, x_2, x_3) = -4(x_1 + x_2 - x_3)^3 + 8x_2^3 = 0, \end{cases}$$

and hence,

$$\begin{cases} x_1 + x_2 - x_3 = \sqrt[3]{2}x_2 \Rightarrow x_1 - x_3 = (\sqrt[3]{2} - 1)x_2 \\ x_2^3 + 3x_2^2(x_3 - x_1) = 0 \Rightarrow x_2 = 0 \text{ or } x_1 - x_3 = \frac{1}{3}x_2, \end{cases}$$

i.e.,  $x_2 = 0, x_1 = x_3$ . Therefore, the stationary points are  $y = (x_1, 0, x_1)$  on the non-negative orthant, and hence,

$$f(x_1, 0, x_1) = 0 \Rightarrow \mathcal{T}y^4 = 0.$$

So  $\mathcal{T}x^4 \geq 0$  at the stationary points and the boundary points, and then  $\mathcal{T}x^4 \geq 0$  for all  $x \geq 0$ . That is,  $\mathcal{T}$  is copositive.

(2) Let  $t_{1113} = t_{1333} = t_{2333} = t_{1222} = t_{2223} = 1$  and  $t_{1112} = -1$  without loss the generality. Then

$$\mathcal{T}x^4 = (x_1 - x_2 + x_3)^4 + 8(x_1x_2^3 + x_2^3x_3 + x_2x_3^3 - 3x_1x_2^2x_3).$$

Let  $f(x_1, x_2, x_3) = \mathcal{T}x^4$ . Similarly, solving the stationary point equations,

$$\begin{cases} f'_{x_1}(x_1, x_2, x_3) = 4(x_1 - x_2 + x_3)^3 + 8x_2^2(x_2 - 3x_3) = 0, \\ f'_{x_2}(x_1, x_2, x_3) = -4(x_1 - x_2 + x_3)^3 \\ \quad + 8(3x_1x_2^2 + 3x_2^2x_3 + x_3^3 - 6x_1x_2x_3) = 0, \\ f'_{x_3}(x_1, x_2, x_3) = 4(x_1 - x_2 + x_3)^3 + 8x_2(x_2^2 + 3x_3^2 - 3x_1x_2) = 0, \end{cases}$$

that is,

$$\begin{cases} (x_1 - x_2 + x_3)^3 + 2x_2^2(x_2 - 3x_3) = 0 \\ 3x_2(x_3^2 - x_1x_2 + x_2x_3) = 0 \Rightarrow x_2 = 0 \text{ or } x_1x_2 = x_3(x_3 + x_2) \\ x_2^3 + x_3^3 + 3x_1x_2(x_2 - 2x_3) = 0, \end{cases}$$

This yields  $x_1 = x_2 = x_3 = 0$  on the non-negative orthant. Therefore, the stationary point is  $y = (0, 0, 0)$ , and hence,  $\mathcal{T}y^4 = f(0, 0, 0) = 0$ . So  $\mathcal{T}x^4 \geq 0$  at the stationary points and the boundary points, and then  $\mathcal{T}x^4 \geq 0$  for all  $x \geq 0$ . That is,  $\mathcal{T}$  is copositive.  $\square$

Combing the conclusions of Theorems 3.5 and 3.7, the main result is established in this subsection.

**Theorem 3.8.** Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 3-dimensional symmetric tensor with  $|t_{ijkl}| = t_{iiii} = t_{iijj} = 1$  for all  $i, j, k, l \in \{1, 2, 3\}$ . Then  $\mathcal{A}$  is copositive if and only if

- (1) there is at most one  $-1$  in  $\{t_{1123}, t_{1223}, t_{1233}\}$ ;
- (2) there is only two  $-1$  in  $\{t_{1123}, t_{1223}, t_{1233}\}$  and there is at least two 1 in  $\{t_{iijj}; i, j = 1, 2, 3, i \neq j\}$ ;
- (3)  $t_{1123} = t_{1223} = t_{1233} = -1$  and there is at least five 1 in  $\{t_{iijj}; i, j = 1, 2, 3, i \neq j\}$ .

### 3.3 Applications of a gerenal tensor

In this subsection, we apply Theorems 3.4 and 3.8 to find the (strict) copositivity of a gerenal 4th-order 3-dimensional symmetric tensor, and moreover, these analytic conditions can be very easily parsed and verified.

For a 4th-order 3-dimensional symmetric tensor  $\mathcal{T} = (t_{ijkl})$  with its entires  $t_{iiii} > 0$  for all  $i \in \{1, 2, 3\}$ , let  $\mathcal{T}' = (t'_{ijkl})$  be a symmetric tensor with its entires  $t'_{1111} = t'_{2222} = t'_{3333} = 1$  and

$$\begin{aligned} t'_{1112} &= t_{1112}t_{1111}^{-\frac{3}{4}}t_{2222}^{-\frac{1}{4}}, t'_{1122} = t_{1122}t_{1111}^{-\frac{1}{2}}t_{2222}^{-\frac{1}{2}}, t'_{1222} = t_{1222}t_{1111}^{-\frac{1}{4}}t_{2222}^{-\frac{3}{4}}, \\ t'_{1113} &= t_{1113}t_{1111}^{-\frac{3}{4}}t_{3333}^{-\frac{1}{4}}, t'_{1133} = t_{1133}t_{1111}^{-\frac{1}{2}}t_{3333}^{-\frac{1}{2}}, t'_{1333} = t_{1333}t_{1111}^{-\frac{1}{4}}t_{3333}^{-\frac{3}{4}}, \\ t'_{2223} &= t_{2223}t_{2222}^{-\frac{3}{4}}t_{3333}^{-\frac{1}{4}}, t'_{2233} = t_{2233}t_{2222}^{-\frac{1}{2}}t_{3333}^{-\frac{1}{2}}, t'_{2333} = t_{2333}t_{2222}^{-\frac{1}{4}}t_{3333}^{-\frac{3}{4}}, \\ t'_{1123} &= t_{1123}t_{1111}^{-\frac{1}{2}}t_{2222}^{-\frac{1}{4}}t_{3333}^{-\frac{1}{4}}, t'_{1223} = t_{1223}t_{1111}^{-\frac{1}{4}}t_{2222}^{-\frac{1}{2}}t_{3333}^{-\frac{1}{4}}, \\ t'_{1233} &= t_{1233}t_{1111}^{-\frac{1}{4}}t_{2222}^{-\frac{1}{4}}t_{3333}^{-\frac{1}{2}}. \end{aligned}$$

For  $y = (y_1, y_2, y_3)^\top$  and  $x = (x_1, x_2, x_3)^\top = (t_{1111}^{\frac{1}{4}}y_1, t_{2222}^{\frac{1}{4}}y_2, t_{3333}^{\frac{1}{4}}y_3)^\top$ , then

$$\begin{aligned} \mathcal{T}y^4 &= t_{1111}y_1^4 + 4t_{1112}y_1^3y_2 + 6t_{1122}y_1^2y_2^2 + 4t_{1222}y_1y_2^3 + t_{2222}y_2^4 \\ &\quad + 4t_{1113}y_1^3y_3 + 6t_{1133}y_1^2y_3^2 + 4t_{1333}y_1y_3^3 + t_{3333}y_3^4 \\ &\quad + 4t_{2223}y_2^3y_3 + 6t_{2233}y_2^2y_3^2 + 4t_{2333}y_2y_3^3 \\ &\quad + 12t_{1123}y_1^2y_2y_3 + 12t_{1223}y_1y_2^2y_3 + 12t_{1233}y_1y_2y_3^2 \\ &= x_1^4 + 4t'_{1112}x_1^3x_2 + 6t'_{1122}x_1^2x_2^2 + 4t'_{1222}x_1x_2^3 + x_2^4 \\ &\quad + 4t'_{1113}x_1^3x_3 + 6t'_{1133}x_1^2x_3^2 + 4t'_{1333}x_1x_3^3 + x_3^4 \\ &\quad + 4t'_{2223}x_2^3x_3 + 6t'_{2233}x_2^2x_3^2 + 4t'_{2333}x_2x_3^3 \\ &\quad + 12t'_{1123}x_1^2x_2x_3 + 12t'_{1223}x_1x_2^2x_3 + 12t'_{1233}x_1x_2x_3^2 \\ &= \mathcal{T}'x^4. \end{aligned}$$



It is obvious that the copositivity of symmetric tensor  $\mathcal{T} = (t_{ijkl})$  is equivalent to the copositivity of  $\mathcal{T}' = (t'_{ijkl})$ . So, applying Corollaries 3.2 and 3.6 (or Theorems 3.4(1) and 3.8(1)) to establish easily the following conclusions .

**Theorem 3.9.** Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 3-dimensional symmetric tensor with  $t_{iiii} > 0$  for all  $i \in \{1, 2, 3\}$  . Assume that one of the following three conditions holds,

- (a)  $t_{1123} \geq -t_{1111}^{\frac{1}{2}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{4}}, t_{1223} \geq t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{2}} t_{3333}^{\frac{1}{4}}, t_{1233} \geq t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{2}};$
- (b)  $t_{1123} \geq t_{1111}^{\frac{1}{2}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{4}}, t_{1223} \geq -t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{2}} t_{3333}^{\frac{1}{4}}, t_{1233} \geq t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{2}};$
- (c)  $t_{1123} \geq t_{1111}^{\frac{1}{2}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{4}}, t_{1223} \geq t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{2}} t_{3333}^{\frac{1}{4}}, t_{1233} \geq -t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{2}}.$

Then (1)  $\mathcal{T}$  is strictly copositive if for all  $i, j \in \{1, 2, 3\}$  and  $i \neq j$ ,

$$t_{iijj} \geq -\sqrt{t_{iiii}t_{jjjj}}, \quad t_{iiij} \geq t_{iiii}^{\frac{3}{4}} t_{jjjj}^{\frac{1}{4}};$$

(2)  $\mathcal{T}$  is copositive if for all  $i, j \in \{1, 2, 3\}$  and  $i \neq j$ ,

$$t_{iijj} \geq \sqrt{t_{iiii}t_{jjjj}}, \quad t_{iiij} \geq -t_{iiii}^{\frac{3}{4}} t_{jjjj}^{\frac{1}{4}}.$$

From Theorems 3.4 (3) and 3.8 (3), the following conclusions are established easily.

**Theorem 3.10.** Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 3-dimensional symmetric tensor with  $t_{iiii} > 0$  for all  $i \in \{1, 2, 3\}$  . Assume that

$$t_{1123} \geq -t_{1111}^{\frac{1}{2}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{4}}, t_{1223} \geq -t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{2}} t_{3333}^{\frac{1}{4}}, t_{1233} \geq -t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{2}}.$$

Then (1)  $\mathcal{T}$  is strictly copositive if for all  $i, j \in \{1, 2, 3\}$  and  $i \neq j$ ,

$$t_{iijj} \geq \sqrt{t_{iiii}t_{jjjj}}, \quad t_{iiij} \geq t_{iiii}^{\frac{3}{4}} t_{jjjj}^{\frac{1}{4}};$$

(2)  $\mathcal{T}$  is copositive if for all  $i, j \in \{1, 2, 3\}$  and  $i \neq j$ ,

$$t_{iijj} \geq \sqrt{t_{iiii}t_{jjjj}}$$

and there is one  $t_{kkkl} \in \{t_{iiij}; i, j = 1, 2, 3, i \neq j\}$  such that

$$t_{kkkl} \geq -t_{kkkk}^{\frac{3}{4}} t_{llll}^{\frac{1}{4}} \text{ and } t_{iiij} \geq t_{iiii}^{\frac{3}{4}} t_{jjjj}^{\frac{1}{4}} \text{ for all } t_{iiij} \neq t_{kkkl}.$$

From Theorems 3.4 (2) and 3.8 (2), the following conclusions are established easily.

**Theorem 3.11.** Let  $\mathcal{T} = (t_{ijkl})$  be a 4th-order 3-dimensional symmetric tensor with  $t_{iiii} > 0$  for all  $i \in \{1, 2, 3\}$  . Assume that one of the following three conditions holds,

- (a)  $t_{1123} \geq -t_{1111}^{\frac{1}{2}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{4}}, t_{1223} \geq -t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{2}} t_{3333}^{\frac{1}{4}}, t_{1233} \geq t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{2}};$   
(b)  $t_{1123} \geq t_{1111}^{\frac{1}{2}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{4}}, t_{1223} \geq -t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{2}} t_{3333}^{\frac{1}{4}}, t_{1233} \geq -t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{2}};$   
(c)  $t_{1123} \geq -t_{1111}^{\frac{1}{2}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{4}}, t_{1223} \geq t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{2}} t_{3333}^{\frac{1}{4}}, t_{1233} \geq -t_{1111}^{\frac{1}{4}} t_{2222}^{\frac{1}{4}} t_{3333}^{\frac{1}{2}}.$

Then (1)  $\mathcal{T}$  is strictly copositive if for all  $i, j \in \{1, 2, 3\}$  and  $i \neq j$ ,

$$t_{iiij} \geq t_{iiii}^{\frac{3}{4}} t_{jjjj}^{\frac{1}{4}};$$

and there is one  $t_{kkll} \in \{t_{iiij}; i, j = 1, 2, 3, i \neq j\}$  such that

$$t_{kkll} \geq \sqrt{t_{kkkk} t_{llll}} \text{ and } t_{iiij} \geq -\sqrt{t_{iiii} t_{jjjj}} \text{ for all } t_{iiij} \neq t_{kkll}.$$

(2)  $\mathcal{T}$  is copositive if for all  $i, j \in \{1, 2, 3\}$  and  $i \neq j$ ,

$$t_{iiij} \geq \sqrt{t_{iiii} t_{jjjj}}$$

and there is two  $t_{kkkl}, t_{sssr} \in \{t_{iiij}; i, j = 1, 2, 3, i \neq j\}$  such that

$$t_{kkkl} \geq t_{kkkk}^{\frac{3}{4}} t_{llll}^{\frac{1}{4}}, \quad t_{sssr} \geq t_{ssss}^{\frac{3}{4}} t_{rrrr}^{\frac{1}{4}}$$

and for all  $t_{iiij} \in \{t_{iiij}; i, j = 1, 2, 3, i \neq j\} \setminus \{t_{kkkl}, t_{sssr}\},$

$$t_{iiij} \geq -t_{iiii}^{\frac{3}{4}} t_{jjjj}^{\frac{1}{4}}.$$

## 4 Conclusions

For a 4th-order 3-dimensional symmetric tensor with its entries 1 or  $-1$ , the analytic necessary and sufficient conditions are established for strict copositivity and copositivity, respectively. These conditions can be applied to verify (strict) copositivity of a general 4th order 3-dimensional symmetric tensor.

## References

- [1] K. Kannike, Vacuum stability conditions from copositivity criteria. Eur. Phys. J. C, 72(2012), 2093
- [2] K. Kannike, Vacuum stability of a general scalar potential of a few fields. Eur. Phys. J. C, 76(2016), 324; Erratum. Eur. Phys. J. C, 78(2018), 355
- [3] H. Chen, Z.H. Huang, L. Qi, Copositive tensor detection and its applications in physics and hypergraphs. Comput. Optim. Appl. 69(2018), 133-158

- [4] Y. Song, X. Li, Copositivity for a class of fourth order symmetric tensors given by scalar dark matter. *J. Optim Theory Appl.* 195 (2022), 334-346
- [5] Y. Song, Co-positivity of tensors and boundedness-from-below conditions of CP conserving two-Higgs-doublets potential. *Inter. J. Modern Phys. A.* **2023**, 38, 2350164.
- [6] Y. Song, Vacuum stability conditions of the general two-Higgs-doublet potential. *Modern Phys. Lett. A.* 38(28n29)(2023), 2350130.
- [7] J. Nie, Z. Yang, X. Zhang, A complete semidefinite algorithm for detecting copositive matrices and tensors, *SIAM Journal on Optimization*, 28 (2018), 2902-2921.
- [8] C. Wang, H. Chen, Y. Wang, H. Yan, G. Zhou, Copositive tensor optimization problem and its applications to hypergraphs. *J. Indus. Manag. Optim.* doi: 10.3934/jimo.2023109
- [9] H. Chen, Y. Wang. High-order copositive tensors and its applications, *J. Appl. Anal. Comput.* 8(6)(2018) 1863-1885
- [10] L. Qi, Symmetric Nonnegative Tensors and Copositive Tensors. *Linear Algebra Appl.*, 439(2013),228-238
- [11] L. Qi, Eigenvalues of a real supersymmetric tensor. *J. Symbolic Comput.*, 40(6) (2005), 1302-1324
- [12] T.S. Motzkin, Copositive quadratic forms, National Bureau of Standards Report, 1818 (1952), 11-12
- [13] V. Baston, Extreme copositive quadratic forms. *Acta Arithmetica* 15 (1969), 319-327
- [14] H. C. Simpson, S. J. Spector, On copositive matrices and strong ellipticity for isotropic elastic materials, *Arch. Rational Mech. Anal.*, 84 (1983), pp. 55–68.
- [15] K.P. Hadeler, On copositive matrices. *Linear Algebra Appl.* 49(1983), 79-89
- [16] E. Nadler, Nonnegativity of bivariate quadratic functions on a triangle. *Comput. Aided Geom. D.* 9(1992),195-205
- [17] G. Chang, T.W. Sederberg, Nonnegative quadratic Bézier triangular patches. *Comput. Aided Geom. D.* 11(1994), 113-116
- [18] L.E. Andersson, G. Chang, T. Elfving, Criteria for copositive matrices using simplices and barycentric coordinates. *Linear Algebra Appl.* 5(1995), 9-30
- [19] J.W. Schmidt, W. Heß, Positivity of cubic polynomials on intervals and positive spline interpolation, *BIT Numerical Mathematics*, 28(2)(1988), 340-352

- [20] L. Qi, Y. Song, X. Zhang, Positivity Conditions for Cubic, Quartic and Quintic Polynomials. *J. Nonlinear Convex Anal.* 23(2)(2022), 191-213
- [21] J. Liu, Y. Song, Copositivity for 3rd order symmetric tensors and applications. *Bull. Malays. Math. Sci. Soc.* 45(1)(2022), 133-152.
- [22] G. Ulrich, L.T. Watson, Positivity conditions for quartic polynomials. *SIAM J. Sci. Comput.* 15(1994), 528-544
- [23] L. Qi, Y. Song, X. Zhang, Copositivity of Three-Dimensional Symmetric Tensors. *Asia-Pacific J. Opera. Research* 40(2)(2023), 2250032.
- [24] H. Chen, Z. Huang, L. Qi, Copositivity Detection of Tensors: Theory and Algorithm. *J. Optim. Theory Appl.* 174(2017), 746-761
- [25] L. Li, X. Zhang, Z. Huang, L. Qi, Test of copositive tensors. *J. Industrial Manag. Optim.* 15(2018), 881-891
- [26] Y. Song, L. Qi, Necessary and sufficient conditions for copositive tensors. *Linear Multilinear A* 63(1)(2015), 120-131
- [27] Y. Song, L. Qi, Eigenvalue analysis of constrained minimization problem for homogeneous polynomial. *J. Glob. Optim.* 64(2016), 563-575
- [28] Y. Song, L. Qi, Tensor Complementarity Problem and Semi-positive Tensors. *J Optim Theory Appl* 169(2016), 1069-1078
- [29] M. Laurent, L. F. Vargas, Exactness of Parrilo's Conic Approximations for Copositive Matrices and Associated Low Order Bounds for the Stability Number of a Graph. *Math. Oper. Res.* 48(2) (2022) 1017-1043.
- [30] M. Hall and M. Newman, Copositive and completely positive quadratic forms, *Proc. Cam. Philos. Soc.*, 59 (1963), 329-339.
- [31] P. Diananda, On non-negative forms in real variables some or all of which are non-negative, *Math. Proc. Camb. Philos. Soc.* 58 (1) (1962) 17-25
- [32] Leonard D. Baumert, Extreme copositive quadratic forms, *Pacific J. Math.* 19 (1966) 197-204.
- [33] Leonard D. Baumert, Extreme copositive quadratic forms, II, *Pacific J. Math.* 20 (1967) 1-20.
- [34] J.-B. Hiriart-Urruty, A. Seeger, A variational approach to copositive matrices, *SIAM Rev.* 52 (4) (2010) 593-629.
- [35] L. Qi, H. Chen, Y. Chen, *Tensor Eigenvalues and Their Applications*. Springer Singapore, 2018
- [36] L. Qi, L., Z. Luo, *Tensor Analysis: Spectral Theory and Special Tensors*. SIAM, Philadelphia 2017

- [37] M. Dong and H. Chen, Geometry of the Copositive Tensor Cone and its Dual. *Asia-Pacific J. Opera. Res.* 37(4) (2020), 2040008
- [38] Z. Huang, L. Qi, Tensor Complementarity Problems—Part I: Basic Theory. *J Optim Theory Appl* 183(2019), 1–23
- [39] L. Qi, Z. Huang, Tensor Complementarity Problems—Part II: Solution Methods. *J Optim Theory Appl* 183(2019), 365–385
- [40] Z. Huang, L. Qi, Tensor Complementarity Problems—Part III: Applications. *J Optim Theory Appl* 183(2019), 771–791
- [41] R. Meng, Z. Huang, Y. Wang, Existence of the least element solution of the vertical block Z-tensor complementarity problem. *Optim Lett* 17(2023), 1697–1709
- [42] S. Sharma, K. Palpandi, A criterion for Q-tensors. *Optim Lett* (2023). doi: 10.1007/s11590-023-02074-w
- [43] X. Li, G. Tang, Strict feasibility for the polynomial complementarity problem. *J Glob Optim* (2023). doi: 10.1007/s10898-023-01339-z
- [44] R. Balaji, K. Palpandi, Positive definite and Gram tensor complementarity problems. *Optim. Lett.* 12(2018), 639–648
- [45] X. Bai, Z. Huang, Y. Wang, Global uniqueness and solvability for tensor complementarity problems. *J. Optim. Theory Appl.* 170(2016), 72–84
- [46] M. Che, L. Qi, L.Y. Wei, Stochastic  $\mathcal{R}_0$  tensors to stochastic tensor complementarity problems. *Optim. Lett.* 13(2019), 261–279
- [47] M. Che, L. Qi, Y. Wei, Positive-definite tensors to nonlinear complementarity problems. *J. Optim. Theory Appl.* 168(2016), 475–487
- [48] H. Chen, Y. Wang, High-order copositive tensors and its applications. *J. Appl. Anal. Compu.* 8(2018), 1863–1885
- [49] H. Chen, L. Qi, Y. Song, Column sufficient tensors and tensor complementarity problems. *Front. Math. China*, 13(2018), 255–276
- [50] M.S. Gowda, Polynomial complementarity problems. *Pac. J. Optim.* 13(2017), 227–241
- [51] Z. Huang, L. Qi, Formulating an  $n$ -person noncooperative game as a tensor complementarity problem. *Comput. Optim. Appl.* 66(2017), 557–576
- [52] Z. Luo, L. Qi, N. Xiu, The sparsest solutions to  $Z$ -tensor complementarity problems. *Optim. Lett.* 11(2017), 471–482
- [53] Y. Song, G. Yu, Properties of solution set of tensor complementarity problem. *J. Optim. Theory Appl.* 170(2016), 85–96
- [54] Y. Song, L. Qi, Strictly semi-positive tensors and the boundedness of tensor complementarity problems. *Optim. Lett.* 11(2017), 1407–1426

- [55] Y. Song, L. Qi, Properties of some classes of structured tensors. J Optim. Theory Appl. 165(2015), 854-873
- [56] Y. Song, W. Mei, Structural Properties of Tensors and Complementarity Problems. J. Optim. Theory Appl. 176(2018), 289-305
- [57] X. Wang, H. Chen, Y. Wang, Solution structures of tensor complementarity problem. Front. Math. China, 13(2018), 935-945
- [58] Y. Wang, Z. Huang, X. Bai, Exceptionally regular tensors and tensor complementarity problems. Optim. Method. Softw. 31(2016), 815-828
- [59] J. Wang, S., Hu, Z. Huang, Solution sets of quadratic complementarity problems. J. Optim. Theory Appl. 176(2018), 120-136